



balkan.bio
2025

6TH BALKAN CONFERENCE ON BIOSCIENCES



BOOK OF ABSTRACTS AND PROGRAM

ISBN 978-619-281-077-1



30-31 OCT 2025
PLOVDIV, BULGARIA



Committees

Organizing Committee:

Prof. Vesselin Baev, University of Plovdiv

Assoc. Prof. Anastas Gospodinov, The Institute of Molecular Biology (IMB, BAS)

Assoc. Prof. Elena Apostolova, University of Plovdiv

Assoc. Prof. Slaveya Petrova, University of Plovdiv

Assoc. Prof. Penka Vasileva, University of Plovdiv

Assoc. Prof. Vesela Mitkovska, University of Plovdiv

Assist. Prof. Vesela Yancheva, University of Plovdiv

Assist. Prof. Gergana Zahmanova, University of Plovdiv

Assist. Prof. Krasimir Todorov, University of Plovdiv

Assist. Prof. Bogdan Nikolov, University of Plovdiv

Assist. Prof. Boryana Georgieva, University of Plovdiv

Dr. Marinela Hadzhieva, University of Plovdiv

Dr. Daniella Mollova, University of Plovdiv

Scientific Committee:

Prof. Velizar Gochev, University of Plovdiv

Prof. Teodora Staykova, University of Plovdiv

Prof. Galina Yahubyan, University of Plovdiv

Prof. Stefan Dimitrov, The Institute of Molecular Biology (IMB, BAS)

Prof. Iva Ugrinova, The Institute of Molecular Biology (IMB, BAS)

Assoc. Prof. Anastas Gospodinov, The Institute of Molecular Biology (IMB, BAS)

Prof. Plamen Stoyanov, University of Plovdiv

Prof. Elenka Georgieva, University of Plovdiv

Prof. Dilyan Georgiev, University of Plovdiv

Prof. Sonya Kostadinova, University of Plovdiv

Prof. Valentina Margaritova, University of Plovdiv

Assoc. Prof. Galina Radeva, The Institute of Molecular Biology (IMB, BAS)

Assoc. Prof. Dimitar Iliev, The Institute of Molecular Biology (IMB, BAS)

Assoc. Prof. Emilia Andreenko, University of Plovdiv

Assoc. Prof. Hristo Dimitrov, University of Plovdiv

P162 Evaluating the use of saline diatom indices for bioassessment through metabarcoding and microscopy in two inland saline lakes (Serbia)

Ljubica Vlaović¹, Clarisse Lemonnier², Benjamin Alric^{2,3}, Frédéric Rimet^{2,3}, Miloš Ćirić⁴, Srđan Miletić⁴, Željka Milovanović⁴, Vladimir Petrović⁴, Jelena Krizmanić¹, Danijela Vidaković⁴

¹*University of Belgrade, Faculty of Biology, Institute of Botany and Botanical Garden
“Jevremovac“, Takovska 43, Belgrade 11000, Serbia*

²*Université Savoie-Mont Blanc, INRAE, UMR CARTELE, 74200 Thonon-les-Bains, France*

³*Pôle R&D ECLA, INRAE, 74200 Thonon-les-Bains, France*

⁴*University of Belgrade, Institute of Chemistry, Technology and Metallurgy, National Institute of the Republic of Serbia, Njegoševa 12, Belgrade 11000, Serbia*

e-mail: ljubica.vlaovic2507@gmail.com

Keywords: diatoms, saline lakes, metabarcoding, microscopy, biomonitoring

Diatoms are widely recognized as effective bioindicators of ecological status in aquatic ecosystems, routinely used in biomonitoring. Traditional microscopy-based identification is time-consuming and requires taxonomic expertise, prompting the use of DNA metabarcoding as a faster and complementary or alternative method for bioassessment. This study evaluates the applicability of *rbcL*-based metabarcoding alongside microscopy for the assessment of two Serbian saline lakes—Plava Banja and Pečena Slatina—located in the Pannonian Plain. These lakes host specialized diatom communities dominated by halophilic and halotolerant taxa. To assess their ecological status/potential, two salinity-specific diatom indices were applied: the Diatom Index for Soda Pans (DISP) and the Trait-Based Index (TBI). In Pečena Slatina, both indices yielded comparable values between microscopy and metabarcoding, suggesting methodological consistency. In contrast, results from Plava Banja showed only partial agreement. In both lakes, index values were positioned near the upper end of the scale, indicating good ecological status/potential. However, microscopy consistently included a significantly higher proportion of taxa in index calculations—exceeding 79% in Plava Banja and 90% in Pečena Slatina—compared to 51–65% for metabarcoding. This discrepancy reflects current limitations in reference libraries for saline environments and incomplete trait annotation in molecular datasets, particularly the lack of species-level resolution and ecological or biovolume data. These constraints reduce the robustness of index-based assessments using metabarcoding. While metabarcoding shows promise, its application in saline ecosystems remains limited until gaps in reference libraries and trait coverage are addressed.

Acknowledgments: This work has received funding from the European Union’s Horizon Europe research and innovation programme under grant agreement No 101079234 (BIOLAWEB project) and Alexander von Humboldt Foundation, and was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Grants No. 451-03-136/2025-03/200178, No. 451-03-137/2025-03/200178 and No. 451-03-1041/2024-03/2/3322).